

Hendrich II Fall Risk Model® Validation

Interpreting Studies of Fall Risk Tools

The diagnostic validity of a clinical tool is the extent to which the tool measures what it is designed to measure. With regard to fall risk tools, validity is the extent to which the tool correctly identifies patients at high risk for falls. A number of statistical measures are commonly used to assess the diagnostic validity of fall risk tools, including sensitivity and specificity. Because it is difficult, if not impossible, to create a tool that is 100% valid, diagnostic validity is measured by degrees. For most measures of tool validity, values close to 100% (or to 1, depending on the variable) reflect the optimal state of validity, but it is not realistic in the real world.

An effective fall risk assessment tool for use in the clinical setting should be sensitive enough to identify high-risk persons/patients and specific enough to identify those persons who are not at risk, thereby allowing for the targeted use of evidence-based interventions to minimize injurious falls. A tool should also be simple to use as holistic screening, easy to score, and quick to complete to minimize burden on care provider time. It should also support specific interventions aimed at reducing the underlying cause of the identified risk factors. For example, confusion is not a diagnosis, and yet etiologies of confusion are among the most underdiagnosed factors in many acute care facilities. Ideally, the presence of a risk factor like confusion should trigger an immediate evaluation of the patient; for patients with confusion, this evaluation should differentiate delirium from dementia or medication side effects, for example, and determine whether a more in-depth assessment is needed. In other words, reframing fall risk factors as a window into healthier aging and functionality, while reducing injurious falls, is the real opportunity of using a validated tool in today's care continuum.

Aspects of Study Design Can Affect Validity of Fall Risk Tools

Considerations when interpreting studies of fall risk tools include the study design (e.g., prospective vs. retrospective, case-control vs. uncontrolled, observational vs. interventional, etc.), patient population (e.g., general acute care vs. specific populations), sample size (e.g., large vs. small, number of fall events), duration (long vs. short), and methods of data retrieval and analysis (e.g., chart review vs. prospective data collection). Larger and more diverse samples, particularly of the study event (i.e., falls), may increase the reliability of findings and, possibly, foster broader applicability to real-world populations. For example, the initial validation study of the HIIIFRM¹ enrolled more than 1100 general acute-care patients and recorded 254 fall events over 2 years. This large study included the diverse patient population of a

Level I Trauma center, with skilled nursing and rehabilitation onsite (increasing the generalizability of the findings), a large number of events (increasing the reliability of the findings), and a long duration (increasing capture of rare events).

In prospective studies, investigators determine before the study what variables will be assessed, which subjects will be enrolled, how study tools will be administered, the length and sample size of the study, and other parameters. This type of design fosters consistency of tool application and data acquisition. However, prospective studies in which providers administer the risk assessment but do not perform any intervention may suffer from the so-called Hawthorn effect.² This term describes changes in care delivered by providers based on their awareness of risk. In other words, providers who identify patients who are high risk for falls will alter their approach to these patients, possibly reducing the incidence of falls. In contrast, retrospective studies collect data from existing sources (e.g., Electronic Medical Records) and therefore cannot ensure consistency in how the tool is used, how the data are collected, and how subjects are enrolled, often with variability in assessments between providers.

Other aspects of study design that can affect measures of diagnostic validity include the use of fall prevention programs at the study sites. Effective fall reduction programs will skew validity analysis by preventing the outcome of interest (i.e., falls), unless statistical analysis is performed to adjust for this confounder. The study population can also affect the evaluation of validity. An older or sicker population, for example, may have higher fall risk overall, possibly altering measures of tool accuracy and affecting the generalizability of the findings. Indeed, readers of the falls literature must use caution when interpreting the results of studies that incorporate falls prevention programs, specific patient populations (e.g., elderly, pediatric), or settings (e.g., rehabilitation, orthopedics, neurology) that may affect the overall incidence of falls.

Application of the risk assessment tools by clinicians during the studies may differ as well. For example, it is recommended that HIIIFRM be administered repeatedly, including on admission, following changes in patient status, and after a fall event. However, many studies require assessment only on admission, with no re-evaluations during hospital stay, even following surgery or other changes in patient status. This use of the HIIIFRM differs from recommended practice and may miss changes in fall risk, thereby introducing inaccuracies to study findings.

Studies of the HIIIFRM: High Validity Across Populations

The HIIIFRM has been evaluated in multiple studies from the United States, Italy, Portugal, Singapore, China, Korea, and other countries (Table 1).^{1,3-16} Study methodologies vary, but include case-control and prospective and retrospective designs. Validity metrics reported in these studies are relatively consistent: sensitivity in the range of 64.9% to 93.2% and specificity 51.3%

to 89.3%, with positive predictive values (PPV) from 0.95% to 7.5% and negative predictive values (NPV) from 96% to 99.7%. These values reflect the percentage of fallers (PPV) and non-fallers (NPV) correctly identified by the tool.

It should be noted that lower sensitivities and specificities have been reported in some studies. For example, an Italian study reported a sensitivity of 45.8%, and two groups reported low specificities (43% and 35%), likely a result of the elderly patient populations enrolled in these studies.⁴⁻⁶ Two emergency department (ED) studies reported sensitivities of only 23.8% to 37.5%. In one of these studies, the HIIIFRM was not administered during the study, but rather, assessed only through review of risk factors in patient charts, which would not allow for an accurate functional test.¹³ In the second study, only fall-related visits to the ED were counted as falls, and falls that occurred in the home that did not lead to an ED visit were not counted.¹⁶

References

1. Hendrich AL, Bender PS, Nyhuis A. Validation of the Hendrich II Fall Risk Model: a large concurrent case/control study of hospitalized patients. *Appl Nurs Res.* Feb 2003;16(1):9-21.
2. Streiner D, Norman G, Blum H. *PDQ Epidemiology.* Toronto: Decker BC; 1989.
3. Hendrich A, Nyhuis A, Kippenbrock T, Soja ME. Hospital falls: development of a predictive model for clinical practice. *Appl Nurs Res.* Aug 1995;8(3):129-139.
4. Lovallo C, Rolandi S, Rossetti AM, Lusignani M. Accidental falls in hospital inpatients: evaluation of sensitivity and specificity of two risk assessment tools. *J Adv Nurs.* Mar 2010;66(3):690-696.
5. Ivziku D, Matarese M, Pedone C. Predictive validity of the Hendrich fall risk model II in an acute geriatric unit. *Int J Nurs Stud.* Apr 2011;48(4):468-474.
6. Caldevilla MN, Costa MA, Teles P, Ferreira PM. Evaluation and cross-cultural adaptation of the Hendrich II Fall Risk Model to Portuguese. *Scand J Caring Sci.* Jun 2013;27(2):468-474.
7. Chapman J, Bachand D, Hyrkas K. Testing the sensitivity, specificity and feasibility of four falls risk assessment tools in a clinical setting. *J Nurs Manag.* Jan 2011;19(1):133-142.
8. Kim EA, Mordiffi SZ, Bee WH, Devi K, Evans D. Evaluation of three fall-risk assessment tools in an acute care setting. *J Adv Nurs.* Nov 2007;60(4):427-435.
9. Nassar N, Helou N, Madi C. Predicting falls using two instruments (the Hendrich Fall Risk Model and the Morse Fall Scale) in an acute care setting in Lebanon. *J Clin Nurs.* Oct 11 2013.
10. Yip WK, Mordiffi SZ, Wong HC, Ang EN. Development and Validation of a Simplified Falls Assessment Tool in an Acute Care Setting. *J Nurs Care Qual.* Oct-Dec 2016;31(4):310-317.
11. Jung H, Park HA. Testing the Predictive Validity of the Hendrich II Fall Risk Model. *West J Nurs Res.* Dec 2018;40(12):1785-1799.
12. Zhang C, Wu X, Lin S, Jia Z, Cao J. Evaluation of Reliability and Validity of the Hendrich II Fall Risk Model in a Chinese Hospital Population. *PLoS One.* 2015;10(11):e0142395.
13. Terrell KM, Weaver CS, Giles BK, Ross MJ. ED patient falls and resulting injuries. *J Emerg Nurs.* Apr 2009;35(2):89-92.
14. Van Dyke D, Singley B, Speroni KG, Daniel MG. Evaluation of fall risk assessment tools for psychiatric patient fall prevention: a comparative study. *J Psychosoc Nurs Ment Health Serv.* Dec 1 2014;52(12):30-35.
15. Campanini I, Mastrangelo S, Bargellini A, et al. Feasibility and predictive performance of the Hendrich Fall Risk Model II in a rehabilitation department: a prospective study. *BMC Health Serv Res.* Jan 11 2018;18(1):18.
16. Patterson BW, Repplinger MD, Pulia MS, et al. Using the Hendrich II Inpatient Fall Risk Screen to Predict Outpatient Falls After Emergency Department Visits. *J Am Geriatr Soc.* Apr 2018;66(4):760-765.

Summary

The HIIIFRM compares very favorably to other tools in terms of sensitivity, specificity, and positive and negative predictive value in multiple clinical studies and diverse patient populations. Moreover, the HIIIFRM is the only tool to incorporate independent risk factors identified through systematic evaluation and regression analysis of more than 600 variables reported in peer-reviewed literature. These qualities make the HIIIFRM a truly predictive tool and allow for the targeting of interventions to specific categories of injurious fall risk with a larger opportunity to build a care continuum approach that can reduce modifiable risk factors whenever possible to preserve health and function. Removing the underlying cause of risk can alter a person's risk state if the provider views risk factors as a holistic approach to care and management.

Hendrich II Fall Risk Model® Validation

Table 1. Diagnostic validation studies of the Hendrich II Fall Risk Model (HIIFRM)

Study	N	Setting	Study design	Sensitivity, %	Specificity, %	PPV/NPV, %	AUC	Notes
General Acute Care Inpatients								
Hendrich et al, 1995 ^{1*}	102 falls 236 control	General acute care	Case-control	77	72	NR	NR	Cross validation with a 1987 data set showed a sensitivity of 83% and specificity of 66%
Hendrich et al, 2003 ²	355 falls 780 controls	General acute care	Case-control	74.9	73.9	NR	NR	HIIFRM validation study
Kim et al, 2007 ³	5489 total	Acute care	Descriptive, prospective	70	61.5	2/99.5	0.73	Comparative study at single hospital in Singapore
Lovallo et al, 2010 ⁴	1148 total 59 falls	Acute care, patients > 50 years	Prospective, observational	45.8	71.0	6/96	NR	Comparative study at medical, surgical, and rehabilitation units at an Italian hospital
Chapman et al, 2011 ⁵	1540 total	Acute care	Descriptive, cross-sectional	64.9	69	7.5/98.1	NR	Comparative study of four fall-risk assessment tools in 17 units
Nassar et al, 2014 ⁶	1815 total	Acute care	Prospective	55.2	89.3	16.5/98.3	NR	Comparative study in Lebanese acute care hospital
Kim et al, 2013 ⁷	1026 total 32 falls	Acute care (neurological patients)	Prospective, descriptive	59.4	78.5	8.2/98.4	0.74	Comparative study of neurology, neurosurgery, and rehabilitation patients at a Korean hospital
Yip et al, 2016 ⁸	10381 total 64 falls	10 general adult wards, patients > 21 years	Prospective observational	75	51.31	.95/99.7	0.67	Comparative study at single hospital in Singapore
Jung and Park, 2018 ⁹	15,170 controls 310 falls	Acute care	Retrospective case-control	67.4-80.0	59.5-64.0	4/99	0.70-0.74	Evaluated on neurology, neurosurgery, hematology, and oncology units (255 beds) in tertiary care hospital in Korea

Hendrich II Fall Risk Model[®] Validation

Study	N	Setting	Study design	Sensitivity, %	Specificity, %	PPV/NPV, %	AUC	Notes
General Acute Care Inpatients								
Cho et al, 2018 ¹⁰	14,307 total 238 falls	Acute care	Retrospective	NR	NR	NR	0.69	Modeling study conducted at tertiary care hospital in Korea
Cho et al, 2020 ¹¹	447 falls 1,341 controls	Acute care	Retrospective, case-control	57.27	78	46.5/84.6	74.2	Comparative study at tertiary care hospital in Korea
Hendrich et al, 2020 ¹²	625 falls 213,733 non-falls	Acute care and critical access	Retrospective	78.72	64.07	0.64/99.9	0.765	Nine hospitals with patient populations representative of US diversity of race, gender, age, ethnicity, case-mix, length of stay, licensed bed size, academic and nonacademic settings, and urban and rural locations. All levels and acuity of inpatient care, including behavioral health, skilled nursing, observation, and ED
Geriatric Inpatients (> 60 years of age)								
Ivziku et al, 2011 ¹³	179 total 14 falls	Geriatric acute care, patients > 65 years	Descriptive, prospective	86	43	11/97	0.72	Italian validation study
Caldevilla et al, 2013 ¹⁴	586 total 104 falls	Acute care, patients > 65 years	Prospective	93.2	35	17.2/97.3	0.65	Portuguese validation study
Zhang et al, 2015 ¹⁵	989 total 32 falls	Acute care, patients > 60 years with chronic diseases	Prospective cross-sectional	72.0	69.0	7/98	0.82	Chinese validation study

Hendrich II Fall Risk Model® Validation

Study	N	Setting	Study design	Sensitivity, %	Specificity, %	PPV/NPV, %	AUC	Notes
Other Clinical Settings								
Terrell et al, 2009 ¹⁶	57 falls	Emergency Department	Retrospective chart review	37.5	NR	NR	NR	HIIFRM was not administered during study; risk factors estimated via chart review
Van Dyke et al, 2014 ¹⁷	50 total 2 falls	12-bed rural psychiatric unit	Prospective	100	67.8	1.9/100	NR	Pilot study comparing two tools in adult psychiatric unit
Campanini et al, 2018 ¹⁸	147 total 11 falls	Inpatient Rehabilitation	Prospective	100	49	NR	0.78	Italian rehabilitation department with patients from orthopedic, pulmonary, and neurology services; HIIFRM administered only on admission to unit
Baran and Gunes, 2018 ¹⁹	159 total 51 falls	Nursing home, patients >65 years	Prospective, observational	80.4	43.5	40.2/82.5	0.62	Comparative study conducted a nursing home in Turkey
Patterson et al, 2018 ²⁰	4366 total 363 falls with ED visit	Emergency Department, patients > 65 years	Retrospective chart review	23.8	90.0	NR	0.64	Outcome was fall-related visit to ED within 6 months of index ED visit, population limited to discharged patients > 65 years of age with PCP in network

*Previous version of HIIFRM, which did not include Get-Up-and-Go test.

AUC: area under the receiver operating characteristic curve; PPV/NPV: positive predictive value/negative predictive value; NR: not reported; ED: emergency department; PCP: primary care physician

References

1. Hendrich A, Nyhuis A, Kippenbrock T, Soja ME. Hospital falls: development of a predictive model for clinical practice. *Appl Nurs Res*. Aug 1995;8(3):129-139.
2. Hendrich AL, Bender PS, Nyhuis A. Validation of the Hendrich II Fall Risk Model: a large concurrent case/control study of hospitalized patients. *Appl Nurs Res*. Feb 2003;16(1):9-21.
3. Kim EA, Mordiffi SZ, Bee WH, Devi K, Evans D. Evaluation of three fall-risk assessment tools in an acute care setting. *J Adv Nurs*. Nov 2007;60(4):427-435.
4. Lovallo C, Rolandi S, Rossetti AM, Lusignani M. Accidental falls in hospital inpatients: evaluation of sensitivity and specificity of two risk assessment tools. *J Adv Nurs*. Mar 2010;66(3):690-696.
5. Chapman J, Bachand D, Hyrkas K. Testing the sensitivity, specificity and feasibility of four falls risk assessment tools in a clinical setting. *J Nurs Manag*. Jan 2011;19(1):133-142.
6. Nassar N, Helou N, Madi C. Predicting falls using two instruments (the Hendrich Fall Risk Model and the Morse Fall Scale) in an acute care setting in Lebanon. *J Clin Nurs*. Oct 11 2013.
7. Kim SR, Yoo SH, Shin YS, et al. Comparison of the reliability and validity of fall risk assessment tools in patients with acute neurological disorders. *Korean Journal of Adult Nursing*. 2013;25(1):24-32.
8. Yip WK, Mordiffi SZ, Wong HC, Ang EN. Development and Validation of a Simplified Falls Assessment Tool in an Acute Care Setting. *J Nurs Care Qual*. Oct-Dec 2016;31(4):310-317.
9. Jung H, Park HA. Testing the Predictive Validity of the Hendrich II Fall Risk Model. *West J Nurs Res*. Dec 2018;40(12):1785-1799.
10. Cho I, Boo EH, Chung E, Bates DW, Dykes P. Novel approach to inpatient fall risk prediction and its cross-site validation using time-variant data. *Journal of Medical Internet Research*. 2019;21(2):e11505.
11. Cho EH, Woo YJ, Han A, Chung YC, Kim YH, Park HA. Comparison of the predictive validity of three fall risk assessment tools and analysis of fall-risk factors at a tertiary teaching hospital. *J Clin Nurs*. 2020;29(17-18):3482-3493. doi:10.1111/jocn.15387
12. Hendrich AL, Bufalino A, Groves C. Validation of the Hendrich II Fall Risk Model: The imperative to reduce modifiable risk factors. *Appl Nurs Res*. 2020;53:151243. doi:10.1016/j.apnr.2020.151243
13. Ivziku D, Matarese M, Pedone C. Predictive validity of the Hendrich fall risk model II in an acute geriatric unit. *Int J Nurs Stud*. Apr 2011;48(4):468-474.
14. Caldevilla MN, Costa MA, Teles P, Ferreira PM. Evaluation and cross-cultural adaptation of the Hendrich II Fall Risk Model to Portuguese. *Scand J Caring Sci*. Jun 2013;27(2):468-474.
15. Zhang C, Wu X, Lin S, Jia Z, Cao J. Evaluation of Reliability and Validity of the Hendrich II Fall Risk Model in a Chinese Hospital Population. *PLoS One*. 2015;10(11):e0142395.
16. Terrell KM, Weaver CS, Giles BK, Ross MJ. ED patient falls and resulting injuries. *J Emerg Nurs*. Apr 2009;35(2):89-92.
17. Van Dyke D, Singley B, Speroni KG, Daniel MG. Evaluation of fall risk assessment tools for psychiatric patient fall prevention: a comparative study. *J Psychosoc Nurs Ment Health Serv*. Dec 1 2014;52(12):30-35.
18. Campanini I, Mastrangelo S, Bargellini A, et al. Feasibility and predictive performance of the Hendrich Fall Risk Model II in a rehabilitation department: a prospective study. *BMC Health Serv Res*. Jan 11 2018;18(1):18.
19. Baran L, Gunes U. Predictive validity of three fall risk assessment tools in nursing home residents in Turkey: A comparison of the psychometric properties. *International Journal of Caring Sciences*. 2018;11(1):36-44.
20. Patterson BW, Reppinger MD, Pulia MS, et al. Using the Hendrich II Inpatient Fall Risk Screen to Predict Outpatient Falls After Emergency Department Visits. *J Am Geriatr Soc*. Apr 2018;66(4):760-765.